



OPEN

DATA DESCRIPTOR

ReSurveyGermany: Vegetation-plot time-series over the past hundred years in Germany

Ute Jandt *et al.*[#]

Vegetation-plot resurvey data are a main source of information on terrestrial biodiversity change, with records reaching back more than one century. Although more and more data from re-sampled plots have been published, there is not yet a comprehensive open-access dataset available for analysis. Here, we compiled and harmonised vegetation-plot resurvey data from Germany covering almost 100 years. We show the distribution of the plot data in space, time and across habitat types of the European Nature Information System (EUNIS). In addition, we include metadata on geographic location, plot size and vegetation structure. The data allow temporal biodiversity change to be assessed at the community scale, reaching back further into the past than most comparable data yet available. They also enable tracking changes in the incidence and distribution of individual species across Germany. In summary, the data come at a level of detail that holds promise for broadening our understanding of the mechanisms and drivers behind plant diversity change over the last century.

Background & Summary

The current biodiversity crisis threatens an estimated one million species with extinction¹. The nature and rate of observed changes depend on the spatial scale at which they are observed². At the finest scale, i.e. the local scale of plant communities, vegetation-plot records have been found to become sometimes richer, sometimes poorer in species³, while a considerable temporal species turnover is apparent in the majority of cases⁴.

Germany has a long tradition in resurvey studies as forest inventories were established already in the 19th century⁵. However, these inventories by default only include tree species and provide no information on other growth forms, and thus, on total vascular plant diversity. In contrast, vegetation scientists carried out plot surveys, so-called relevés, since the beginning of the 20th century⁶, and some of these plots have been repeatedly surveyed. Such vegetation-plot time series have mainly been collected for particular habitats, such as forests^{7–19}, hedgerows²⁰, wet grasslands^{21–24}, mesic grasslands^{25–31}, dry grasslands^{24,32–37}, acid grasslands and heathlands^{38–40}, alpine grasslands^{41,42}, rivers⁴³, riverbanks⁴⁴, peatlands^{45–48}, roadsides⁴⁹ or arable land^{50–52}. Sometimes, they were recorded to assess the changes in species composition across all communities that occur in a certain area^{53–57}. So far, vegetation-plot time series have not been accessible without restrictions. In contrast, open access biodiversity time-series data, such as BioTIME⁵⁸, comprise all different types of taxonomic groups, ranging from plants, plankton and terrestrial invertebrates to vertebrates, but include only a few vegetation-plot time series. Thus, our database closes a gap for a particular region, which is Germany.

Vegetation-plot resurvey data have been extensively used to assess biodiversity changes by means of monitoring certain vegetation types in local studies, such as managed grasslands²⁶ and rivers⁴³. More recently, time series have been collected across regions, exploring the contribution of local biodiversity change³ to that observed at broader spatial scales^{1,59,60}. While these analyses often failed to detect changes in species richness^{3,61,62}, they were able to relate the observed trends to changes in land use and climate^{63,64}. Although these studies have compiled databases on vegetation-plot time series, they are currently not openly available. This is also the case for the current initiative of ReSurveyEurope, which collates and mobilizes vegetation-plot data with repeated measurements over time (<http://euroveg.org/eva-database-re-survey-europe>). Our aim is to provide a comprehensive and taxonomically standardised database of vegetation-plot time series for Germany. We confined the geographical extent to Germany because of a long tradition of German vegetation scientists carrying out temporal observations on

[#]A full list of authors and their affiliations appears at the end of the paper.




Fig. 1 Temporal coverage of the 92 projects included in the study. The coloured lines indicate the start and the end of a project, black diamonds show in which years surveys were made. Resurvey type refers to either studies that were repeated within a particular community across a site without attempts to match plots (community comparison), or were carried out on matched plots, which were either permanently marked or retrieved from exact descriptions (semi-permanent). The lower graph shows the number of times a particular year was included in the covered time span of any of the projects. For a list of projects see Supplementary Table S1.

permanent plots (e.g.³⁰), the large amount of available data, our familiarity with the regional literature, and of recent initiatives to mobilize retrospective biodiversity data for trend analyses (www.idiv.de/smon).

Vegetation-plot time series differ in some fundamental ways from other biodiversity time series. Since the advent of phytosociology in the early 20th century^{65,66}, vegetation surveys in Europe were carried out in a standardised way. Plot sizes of vegetation relevés can vary considerably and depend on the vegetation type considered (e.g. forest plots usually have plot sizes between 100 and 1000 m², while non-forest plots mostly range from 4 to 100 m²⁶⁷). In addition, sampling protocols might vary between studies, but they all include complete lists of species occurring at the plot at the time of sampling. In consequence, vegetation-plot records provide information




Fig. 2 Map of all plots of all projects ($n = 23,641$). Note that green dots may represent one or several plots which were summarised under the same plot resurvey ID ($n = 7,738$). The more complete coverage of Bavaria resulted from including the grassland monitoring Bavaria which started in 2002²⁶.

on both presences and absences of species in a community. As sampling is usually done by professionals, absences of a previously occurring species in a time series strongly indicates local extinction, or vice versa, the presence of a species that had not been recorded previously is a robust indication of colonization. However, even with experts carrying out the survey, it is possible that some species may remain undetected in the record because of their phenology or taxonomic uncertainties⁶⁷. Yet, such vegetation-plot data are much more reliable than vegetation surveys at larger scales, such as floristic grid mapping, where false absence data are common^{68,69}. In contrast to time series at broader spatial scales, vegetation-plot time series contain information on species

Predictors	Population density			Road density			Urban cover			Cropland cover			Protected area		
	Estimates	CI	p	Estimates	CI	P	Estimates	CI	p	Estimates	CI	p	Estimates	CI	p
(Intercept)	30547	30128–30965	<0.001	1.59	1.55–1.63	<0.001	0.07	0.06–0.07	<0.001	0.45	0.44–0.45	<0.001	0.01	0.01–0.01	<0.001
type [unsampled]	–6686	–7447–5925	<0.001	–0.52	–0.59–0.45	<0.001	–0.03	–0.03–0.03	<0.001	0.01	0.00–0.02	0.029	0.01	0.01–0.01	<0.001
Observations	11226			25303			12024			12024			29535		
R ² /R ² adjusted	0.026/0.026			0.008/0.008			0.019/0.019			0.000/0.000			0.038/0.038		

Table 1. Representativeness of grid cells (“Messtischblattquadrant, MTBQ”, a quadrant of the German ordnance maps, 0°5′ × 0°3′) with time series. The estimates were obtained from linear models comparing samples with unsampled MTBQs with respect to population density, road density, urban cover, cropland cover and protected area.

co-occurrence at scales relevant for direct biotic interactions among individuals⁷⁰. An additional advantage of vegetation-plot records is that they report the relative abundance of species, in the case of vegetation records from Germany, typically assessed as cover values^{67,71}. While species cover is very often estimated directly in per cent of ground covered by each species, there is a long tradition in vegetation science of using cover scales with distinct classes to facilitate cover estimations. There is a variety of cover scales, with different classes preferred by researchers in different countries^{71,72}. The still most frequently used scale in Germany was introduced by Braun-Blanquet⁶. This scale, however, is not only based on cover, but uses the abundance of individual plants as additional criterion for species with a cover of ≤1% (classes r and +), which raises difficulties in numerical analyses^{71,73}. To facilitate the estimation of cover changes in time series, Londo introduced a pure cover scale⁷⁴, which in its original or in simplified form (e.g.⁷⁵) became very popular in permanent plot research. It is common practice that resurvey studies use the same cover scale as in the original resurvey. Nevertheless, for a numerical comparison of changes, cover classes have to be converted into per cent values⁷², for which the Turboveg software introduced standardized transformations using the midpoints of the cover classes⁷⁶. The cover information in vegetation-plot records allows key theories of biogeography to be tested, such as the abundance–range size relationship⁷⁷ or the relationship between local abundance and niche breadth^{78,79}. Most importantly, several vegetation-plot time series precede the onset of any other systematic plant species monitoring programme, for example the monitoring of Natura 2000 sites in Europe, which only started in 2001⁸⁰. This is particularly important because severe biodiversity loss may have already happened in the second half of the 20th century, mainly brought about by shifts in the type and intensity of land use as the consequence of technical progress and societal changes⁸¹. Finally, species-abundance data in plots can be linked to functional information on species⁸⁷, which allows the interpretation of the underlying ecological drivers of the changes observed and the consequences for ecosystem functioning⁸².

Based on the data described here we analysed for the first time the dynamics of losses and gains of plant species⁸³. We showed that the difference in cover changes between decreasing and increasing species results in biodiversity change even if species richness at the plot scale remains unchanged. Two mechanisms are responsible for these changes. First, losses at the plot scale were more evenly distributed among losing species than gains among winning species. Second, gains and losses in cover were concentrated in different species, resulting in a higher number of losers than winners at the spatial scale of Germany. The temporal extent of the data allowed us to demonstrate that most species losses occurred already by the 1960s, affecting mostly species from mires and spring fens, grasslands and arable land. Thus, these data already helped to shed light on the most important mechanisms underlying biodiversity change in the second half of the 20th century.

Methods

ReSurveyGermany is the most comprehensive compilation of repeated long-term vegetation plot records from Germany to date, including published studies as well as surveys from grey literature and nature conservation assessments. A list of all 92 projects included in the database is provided in Supplementary Table S1. A project might comprise one or several studies and focus on one or several vegetation types, but typically carried out the surveys at the same times and with the same methodology. In total, the projects comprise 1,794 vascular plant species recorded in 7,738 vegetation plots. The plots were either marked with poles or magnets (permanent) or recovered from exact descriptions, sketches or marks in high-resolution topographic maps (semi-permanent). The uncertainty of the positions varied among studies, but also within a single study as resurveyed plots might have been marked in the later surveys. If the uncertainty was provided by the author or could be estimated from topographic maps, this information was included in the PRECISION field of the header file of ReSurveyGermany (see below). In addition, there were also studies where plots were not matched in time but a set of plots at a site was compared within another set of plots at the same site in the resurvey (community comparison, Fig. 1). We only considered records with complete lists of vascular plants and information on their relative abundance, which was mostly expressed as percentage cover⁸⁴. A further important criterion for including a study was the existence of vegetation data for at least two points in time, although the number of visits (i.e. vegetation records) per site ranged between two and 54. The time span covered by each project is shown in Fig. 1. All records were made between 1927 and 2020. In total, ReSurvey Germany comprises 23,641 vegetation-plot records and 458,311 species cover records.




Fig. 3 Map of plot visits by decade, with the year showing the beginning of the decade.

Plot locations are not evenly spread across Germany (Fig. 2). We assigned the individual plot locations to the grid cells of the quadrants of German ordnance maps (“MTBQ,” $0^{\circ}5' \times 0^{\circ}3'$, approximately $5.6 \text{ km} \times 5.9 \text{ km}$ in the centre of Germany), and tested whether the grid cells with vegetation-plot time-series records differed from those without observations with respect to population density, road density, urban cover, cropland cover and protected areas. Using the land cover dataset from the European Space Agency Climate Change Initiative⁸⁵, we calculated the proportional cover of urban cover for each MTBQ. Spatial information on protected areas was obtained from GIS shapefiles provided by the German Federal Agency for Nature Conservation (Bundesamt für Naturschutz, BfN). This analysis revealed that the sampled grid cells were not representative for the whole area of Germany. As expected from other studies⁸⁶, the sampled grid cells showed significantly higher human

Field name	Type	Description
PROJECT_ID	I	Number of the resurvey project in ReSurveyGermany; see Supplementary Table S1
RELEVE_NR	I	Plot observation ID, only unique within a RS_PROJECT, usually the original plot observation ID from the underlying Turboveg 2 database
PROJECT_ID_RELEVE_NR	C	Unique Plot observation ID, by which the project's plot-species-abundance file is linked to the header file, combination of PROJECT_ID and RELEVE_NR
LAYER	I	0: No layer, 1: Tree layer (uppermost), 2: Tree layer (middle), 3: Tree layer (lowest), 4: Shrub layer (uppermost), 5: Shrub layer (low), 6: Herb layer, 7: Juveniles, 8: Seedling (<1 year), 9: Moss layer.
TaxonName	C	Harmonized taxon name
Cover_Perc	N	Cover of the taxon in per cent

Table 2. Data structure of the Plot-species-abundance file of ReSurveyGermany. For Type: C = character, N = numeric, I = integer (n = 23,641).

population densities, road densities and urban cover, while cover of cropland and the amount of protected area was lower (Table 1), which indicates that the majority of time series was made in regions with higher human pressures. The lack of spatial representativeness also becomes obvious when plotting maps of plot locations by the decade of the times when they were visited (Fig. 3).

While we did not deliberately exclude certain habitat types, the data mainly consist of semi-natural to intensively managed grasslands and forests. Thus, the time series in ReSurveyGermany are biased with respect to habitat types. We assigned EUNIS habitat types to each plot record. The European Nature Information System (EUNIS) provides a comprehensive typology for the terrestrial and marine habitats of Europe⁸⁷. Habitat types are arranged in a hierarchy, from the highest level 1 to the lowest level 4. Here, we show the assignment of plot records to level 3, which was accomplished by using the expert system EUNIS-ESy⁸⁸ and the corresponding R code⁸⁹. Plot records covered a total of 92 EUNIS habitat types out of the 150 ones distinguished for Germany. About 63% of the 23,641 plot records came from grasslands (level 1 EUNIS habitat R, n = 14,849), followed by forests and other wooded lands (T, n = 5,440, 23%). In contrast, arable land (V, vegetated man-made habitats), which makes up more than 36% of the land cover in Germany, was only represented by 3% (816 plot records).

Data Records

The data of the ReSurveyGermany dataset as described above is available <https://doi.org/10.25829/ivid.3514-0qsq70> under the terms specified by CC BY 4.0⁹⁰.

A separate database was created for each project that contributed data, using the data-management software Turboveg 2⁷⁶. The database is composed of two main tables, following the structure of Turboveg and common practice in vegetation science. The plot-species-abundance table contains six fields as described in Table 2. It is linked to the plot metadata (header file) through PROJECT_ID_RELEVE_NR, which is a unique Plot observation ID of a combination of PROJECT_ID (see Supplementary Table S1) and the plot observation ID (called RELEVE_NR), the name of the observed taxon (TaxonName), the vertical layer (tree layer, shrub layer, herb layer, moss layer) in which the species was observed (LAYER) and the taxon's cover in the plot (Cover_Perc). The latter was obtained by transforming the original cover classes in per cent cover, using the midpoints of the cover classes as provided by the Turboveg software⁷⁶. For example, the seven cover classes of the Braun-Blanquet scale⁶, r, +, 1, 2, 3, 4, 5 were transformed to 1%, 2%, 3%, 13%, 38%, 63%, 88%, respectively. The other table is the so-called header file, which holds all important plot-level information, such as plot sizes, geographic location and vegetation structure for each plot observation ID (Table 3).

The taxon names in the plot-species-abundance table were standardised using German SL 1.3⁹¹. The nomenclature for vascular plants followed Wisskirchen *et al.*⁹², with additional aggregations to higher taxonomic levels according to German SL 1.3. As some authors recorded subspecies and other infraspecific taxa, species were aggregated at the species level, using the R package vegdata⁹³. Some closely related species that, from our experience, are often mistaken in the field were merged at the aggregate or genus level. Species aggregates were also used when different taxon names of the same aggregate occurred in different projects, to prevent that the same taxon might appear under different taxon names. We used our own R code to merge taxon names and the notation of the ESy expert system⁸⁸ to protocol all steps. The species harmonisation forms the first section of the ESy system and shows which taxon names were aggregated under the name of a broader taxonomic concept (Supplementary Table S2). In addition, within single projects, we used customised aggregations and segregations when the same taxa were reported with different taxonomic levels at different points in time in the same plot resurvey IDs (Supplementary Table S3). For example, in all years of a time series of a specific plot *Orchis militaris* was reported but in one year *Orchis spec.* was recorded at the genus level. Unaccounted for, such a leap between taxonomic levels within a time series would result in incorrect species change observations. To avoid losing the predominating information at the species level by aggregating all records to *Orchis*, we assumed that the taxon was also *Orchis militaris* in the particular year when only the genus level was reported. If more than one taxon occurred in previous years, we equally distributed the cover values among those taxa. This happened for example when a record was taken late in spring when the two species *Anemone nemorosa* and *A. ranunculoides* could no longer be distinguished.

The percentage cover values of the same aggregated taxon name of the same plot were merged, assuming a random overlap of their cover values and making sure that the combined cover values cannot exceed 100%^{76,94}. This often resulted in cover values with decimal points and might suggest an accuracy of cover estimation that




Fig. 4 Histogram of plot size across all records ($n = 23,641$). Colours show Eunis level 1 habitat types.

is not warranted by the original estimates. As not all projects had recorded cryptogams, we removed bryophytes and lichens in all projects, using the `vegdata` package in R⁹³. As a result, the original list of 3,280 taxon names that included bryophytes and lichens was reduced to 1,794 taxon names of vascular plants. However, if data on lichens and bryophytes are required, they are available on request from the respective dataset custodians (see Supplementary Table S1).

The data structure of the header file of ReSurveyGermany follows the Turboveg 2 standard⁷⁶ and in addition holds the fields of ReSurveyEurope (<http://euroveg.org/eva-database-re-survey-europe>) (Table 3). The fields relevant for the resurvey are `RS_PROJECT`, which refers to the resurvey project in Supplementary Table S1. The header field `RS_SITE` holds the location name of plots and allows for a local geographical scale aggregation of resurvey plots within projects. `LOCALITY` provides more details on the locality in German.

Within each project, the field `RS_PLOT` holds a plot resurvey ID that connects plot observations from different times made on the same plot. In resurveys, there are also cases, where the previously provided location was not precise enough. In these cases, resurveys often used several plots to match one previous plot, resulting in a one-to-many relationship. If a set of plots at the same site was compared with plot records from another point in time, several plot records in the same year might have the same `RS_PLOT` code. The unique code of the one-time observation is a combination of `RS_SITE`, `RS_PLOT` and `YEAR` when the plot was surveyed (`RS_OBSERV`). We report the exact `DATE` when a record was made (if available). In addition, the field `YEAR` lists the year in which the plot was (re)surveyed. If available, we also report the year of the underlying publication (`YEAR_PUBL`).

Plot area (`SURF_AREA`) ranges from 0.5 to 2500 m², with 25, 100 and 400 m² being the most frequently used plot sizes (Fig. 4). Plot sizes larger than 100 m² were typical of forest sites (with a very few exceptions).

Geographic information is given by `LONGITUDE`, `LATITUDE` and `ALTITUDE`. Current monitoring programs and data protection of land owners do not allow us to provide location information at the highest available precision. In addition, some records contain occurrence data of rare and protected species. Thus, information on longitude and latitude was rounded to two decimal digits. Compared to the coordinates at highest available precision, rounding resulted in a mean uncertainty of 371 m (± 138 m standard deviation), and thus, is within the somewhat limited range of accuracy provided by many custodians in the first place (see field `PRECISION`). If more precise coordinates are required for certain analysis we recommend to contact the respective data owners (as shown in Supplementary Table S1). Vegetation-plot time series differ with respect to the accuracy of the plot relocation during the resurvey. In the ideal case, plots are permanently marked, using poles, metal tent pegs or magnets and metal detectors to retrieve their position (shown as “01” in the `LOC_METHOD` field, Table 3). In other cases, plots only have exact coordinates (using GPS coordinates, “03” or “04”) or other ways of descriptions of the exact locality (such as from maps, “05”), but are not marked on the ground, which we refer to as semi-permanent plots. In addition, there is information on the cover scale used for the record, a reference to the data source (or, if published, the publication ID), including the table and column from which the data were taken.

The orientation of the plot can be taken from `SLOPE` (inclination) and slope `ASPECT` (compass directions). Vegetation structure is described by the height and cover of the different layers, ranging from tree layer to moss layer and including information on cover of litter and bare soil (if available).

Some of our projects included experimental treatments with different management of habitats (e.g. abandonment or establishment of grazing, succession and disturbance). Plots with experimental manipulation contain “Y” in the `MANIPULATE` field. The type of manipulation can be taken from `MANIPTYPE`. When projects involved treatments that are not appropriate to assess biodiversity change, we included only the control plots⁴⁶, plots that reflected the predominant land use at the site (e.g. mowing for a grassland to counteract natural succession)²², that were unfenced⁹⁵ or were subjected to continuous grazing⁹⁶.

Field name	Type	Description	Number of NAs
RS_PROJECT	C	Unique name of the resurvey project; for the list of the 92 projects and the underlying references see Supplementary Table S1	0
PROJECT_ID	I	Number of the resurvey project in ReSurveyGermany; see Supplementary Table S1	0
RS_PLOT	C	Unique (within the site) code of the resurveyed plot; it is used to pair observations from different times recorded in the same plot; gives a unique identifier for the resurveyed plot or set of plots in time if combined with RS_PROJECT. Several plots in the same year might have the same RS_PLOT code if they have to be summarised for temporal comparisons. In these cases, they might also contain the community name.	0
RS_SITE	C	Name of the resurveyed site. For further details see LOCALITY.	0
LOCALITY	C	More detailed description of the locality of the resurveyed site, informed by the author (in German and if available)	8,499
RS_OBSERV	C	Unique code of the one-time observation; combination of RS_SITE, RS_PLOT, YEAR	0
RELEVE_NR	I	Plot observation ID, only unique within a RS_PROJECT, usually the original plot observation ID from the underlying Turboveg 2 database	0
PROJECT_ID_RELEVE_NR	C	Unique Plot observation ID, by which the project's plot-species-abundance file is linked to the header file, combination of PROJECT_ID and RELEVE_NR	0
DATE	C	Date of the record (YYYYMMDD); the exact date if provided by the author, otherwise only the year and month or only year; if the year was not provided by the author, we took the year of the publication	0
YEAR	I	Year of the record (YYYY), extracted from DATE	0
SURF_AREA	N	Plot size [m ²] (only stated if available)	2064
MANIPULATE	C	Binary information (Y/N) about whether the plot was part of a manipulative experiment ("Y") or not ("N"). If "Y", we chose the treatments representing the ambient land use. Observations with NA were to our knowledge not part of an experiment, and thus, can be treated as "N".	16,579
MANIPTYPE	C	Shows the type of treatment in the plot manipulation (partly in German and only if available).	20,255
LAND_USE	C	Land use, informed by the authors, often identical with MANIPTYPE (mostly in German, also using the abbreviations used in the particular study, and only if available)	18,149
LOC_METHOD	C	Method of plot (re-)location, 1: Permanently marked plot isolated (i.e. somewhere within the site), 2: Marked plot in a grid (i.e. with regularly spaced neighbor plots), 3: Location with differential GPS, 4: Location with GPS, 5: Location from accurate map, 6: Location from a description, 7: Other	12,607
LOC_METH_COMMENT	C	Detailed description of the location method (if available)	20,163
LONGITUDE	N	Longitude of the plot in decimal degrees, coordinate system WGS-84; this coordinate should refer to the centre of the plot; coordinates were rounded to 2 digits of decimal degrees.	0
LATITUDE	N	Latitude of the plot in decimal degrees, rounded to 2 digits as LONGITUDE	0
PRECISION	I	Uncertainty in m, of coordinates for geographic position of plots, provided by the author or estimated if coordinates were taken from a topographic map. PRECISION refers to the true coordinates, not to those rounded to two digits.	13,034
GEO_LEV	C	Method of how the geographic location was obtained: GPS = Geographical positioning system, MTB = center of the German ordnance map, MTB_4 = center of a quadrant of the German ordnance map, POINT = all other	0
ALTITUDE	I	Elevation [m] (if available)	14,723
ASPECT	N	Compass direction of the slope in degrees [°], 0° = N, 90° = E etc. NA shows plot records either without aspect information or with aspect information when SLOPE is 0. In most cases slope aspect is simply a compass reading and has not been corrected for magnetic declination.	16,572
SLOPE	I	Inclination of the slope in degrees [°]	18,962
COUNTRY	C	DE for Germany	0
EUNIS	C	EUNIS level 3 code of the habitat, as obtained by applying the expert system EUNIS-ESy ⁸⁸ and the corresponding R code ⁸⁹ .	0
COVERSCALE	C	Cover scale used for the plot record. 00 = no scale, cover estimated in per cent (%), 01 = Braun/Blanquet (old), 02 = Braun/Blanquet (new), 03 = Londo, 04 = Presence/Absence, 10 = Reichelt & Wilmanns 1973 (short), 26 = Londo (short), 29 = Londo per cent classes, 30 = Londo (modified, in project 9, Sperle <i>et al.</i> unpublished), 31 = Maas & Kohler 1983 (in project 86), 50 = Londo (modified, in project 89)	0
REFERENCE	C	Reference number in the German Vegetation Reference Database (GVRD), 6 digits referring to the bibliographic reference, found in ReSurveyGermanyReference.csv	0
YEAR_PUBL	I	Year of the publication (if available)	18,057
TABLE_NR	C	Number of the table in the original publication	12,659
TABNAME	C	Name of the table in the original publication	8,402
Continued			

Field name	Type	Description	Number of NAs
NR_IN_TAB	C	Column name in the TABLE_NR	3,789
ORIG_NR	C	Name of the plot given by the author in the original publication	10,172
ORIG_DB	C	Name of original Turboveg file, to be used internally for backtracking changes	19,700
COV_TOTAL	I	Total cover of all layers [%] (if available)	18,704
COV_TREES	I	Cover of the tree layer [%] (if available)	20,520
COV_SHRUBS	I	Cover of the shrub layer [%] (if available)	20,114
COV_HERBS	I	Cover of the herb layer [%] (if available)	11,964
COV_MOSSES	I	Cover of the moss layer (bryophytes and lichens) [%] (if available)	17,512
COV_LITTER	I	Cover of the litter layer on the ground [%] (if available)	20,786
COV_ROCK	I	Cover of the rocks on the plot surface [%] (if available)	21,697
TREE_HIGH	I	Height of the upper tree layer [m] (if available)	22,317
TREE_LOW	I	Height of the lower tree layer [m] (if available)	23,107
SHRUB_HIGH	N	Height of the upper shrub layer [m] (if available)	22,508
SHRUB_LOW	N	Height of the lower shrub layer [m] (if available)	23,478
HERB_HIGH	N	Mean height of the upper herb layer [cm] (if available)	20,311
HERB_LOW	N	Mean height of the lower herb layer [cm] (if available)	22,614
HERB_MAX	N	Maximum height of the herb layer [cm] (if available)	22,627

Table 3. Data structure of the header file of ReSurveyGermany. For Type: C = character, N = numeric, I = integer. Number of NAs = number of missing values out of the total of n = 23,641 records.

Technical Validation

As each dataset was transformed into a Turboveg 2 database⁷⁶, a quality check was made when importing the data. This particularly applied to the taxonomical harmonization of the data, which at the stage of entering the data was adjusted to GermanSL 1.3⁹¹.

Usage Notes

Users are urged to cite the original sources when using ReSurveyGermany in addition to the present paper (see Supplementary Table S1). As some of the time series will be continued, it might be useful to contact the respective data owners. As described above, the dataset cannot be considered representative of Germany's vegetation, neither spatially, nor temporally, which is typical of vegetation-plot time series⁹⁷. As plots were established with different objectives in different habitats at different points in time, analysis of vegetation-plot resurveys faces various methodological challenges⁶². Yet, we note that ReSurveyGermany covers about 60% of the 2,988 vascular plant species that occur in Germany (without subspecies and segregates⁹²) and includes rare habitats which often harbour rare plant species. This means that even if our sites are not fully representative of the vegetation of Germany and its change over the last century, the data nevertheless can provide important insights into biodiversity change at the level of local communities and individual species.

Code availability

The R code to read the plot-species-abundance file (ReSurveyGermany.csv) and combine it with the header data (Header_ReSurveyGermany.csv) is provided on https://github.com/idiv-biodiversity/Read_ReSurveyGermany.

Received: 24 May 2022; Accepted: 7 September 2022;

Published online: 19 October 2022

References

- Díaz, S. *et al.* Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science* **366**, eaax3100 (2019).
- Chase, J. M. *et al.* Species richness change across spatial scales. *Oikos* **128**, 1079–1091 (2019).
- Vellend, M. *et al.* Global meta-analysis reveals no net change in local-scale plant biodiversity over time. *Proceedings of the National Academy of Sciences* **110**, 19456–19459 (2013).
- Blowes, S. A. *et al.* The geography of biodiversity change in marine and terrestrial assemblages. *Science* **366**, 339–345 (2019).
- Riedel, T., Polley, H. & Klatt, S. Germany. in *National Forest Inventories* (eds. Vidal, C., Alberdi, I. A., Hernández Mateo, L. & Redmond, J. J.) 405–421, <https://doi.org/10.1007/978-3-319-44015-6> (Springer International Publishing, 2016).
- Braun-Blanquet, J. *Pflanzensoziologie. Grundzüge der Vegetationskunde*. vol. Seite: (Julius Springer, 1928).
- Bernhardt-Römermann, M. *et al.* Drivers of temporal changes in temperate forest plant diversity vary across spatial scales. *Glob Change Biol* **21**, 3726–3737 (2015).
- Ahrns, C. & Hofmann, G. Vegetationsdynamik und Florenzwandel im ehemaligen mitteldeutschen Waldschutzgebiet 'Hainich' im Intervall 1963–1995. *Hercynia N.F.* **31**, 33–64 (1998).
- Dittmann, T., Heinken, T. & Schmidt, M. Die Wälder von Magdeburgerforth (Fläming, Sachsen-Anhalt) – eine Wiederholungsuntersuchung nach sechs Jahrzehnten, <https://doi.org/10.14471/2018.38.009> (2018).
- Günther, K., Schmidt, M., Quitt, H. & Heinken, T. Veränderungen der Waldvegetation im Elbe-Havelwinkel von 1960 bis 2015. *Tuexenia* **41**, 53–85 (2021).
- Janiesch, P. Vegetationsökologische Untersuchungen in einem Erlenbruchwald im nördlichen Münsterland. 25 Jahre im Vergleich. *Abhandlungen aus dem Westfälischen Museum für Naturkunde* 71–80 (2003).
- Naaf, T. & Wulf, M. Habitat specialists and generalists drive homogenization and differentiation of temperate forest plant communities at the regional scale. *Biological Conservation* **143**, 848–855 (2010).

13. Reinecke, J., Klemm, G. & Heinken, T. Vegetation change and homogenization of species composition in temperate nutrient deficient Scots pine forests after 45 yr. *J Veg Sci* **25**, 113–121 (2014).
14. Mölder, A., Streit, M. & Schmidt, W. When beech strikes back: How strict nature conservation reduces herb-layer diversity and productivity in Central European deciduous forests. *Forest Ecology and Management* **319**, 51–61 (2014).
15. Fischer, C., Parth, A. & Schmidt, W. Vegetationsdynamik in Buchen-Naturwäldern. Ein Vergleich aus Süd-Niedersachsen. *Hercynia N.F.* 45–68 (2009).
16. Schmidt, W. Die Naturschutzgebiete Hainholz und Staufenberg am Harzrand - Sukzessionsforschung in Buchenwäldern ohne Bewirtschaftung (Exkursion E). *Tuexenia* **22**, 151–213 (2002).
17. Strubelt, I., Diekmann, M. & Zacharias, D. Changes in species composition and richness in an alluvial hardwood forest over 52 yrs. *J Veg Sci* **28**, 401–412 (2017).
18. Strubelt, I., Diekmann, M., Peppeler-Lisbach, C., Gerken, A. & Zacharias, D. Vegetation changes in the Hasbruch forest nature reserve (NW Germany) depend on management and habitat type. *Forest Ecology and Management* **444**, 78–88 (2019).
19. Wilmanns, O. & Bogenrieder, A. Veränderungen der Buchenwälder des Kaiserstuhls im Laufe von vier Jahrzehnten und ihre Interpretation - pflanzensoziologische Tabellen als Dokumente. *Abh. Landesmus. Naturk. Münster Westfalen* **48**, 55–79 (1986).
20. Huwer, A. & Wittig, R. Changes in the species composition of hedgerows in the Westphalian Basin over a thirty-five-year period. *Tuexenia* **32**, 31–53 (2012).
21. Immoor, A., Zacharias, D., Müller, J. & Diekmann, M. A re-visitation study (1948–2015) of wet grassland vegetation in the Stedinger Land near Bremen, North-western Germany, <https://doi.org/10.14471/2017.37.013> (2017).
22. Rosenthal, G. Erhaltung und Regeneration von Feuchtwiesen. Vegetationsökologische Untersuchungen auf Dauerflächen. *Diss. Bot.* **182**, 1–283 (1992).
23. Poptcheva, K., Schwartz, P., Vogel, A., Kleinebecker, T. & Hölzel, N. Changes in wet meadow vegetation after 20 years of different management in a field experiment (North-West Germany). *Agriculture, Ecosystems & Environment* **134**, 108–114 (2009).
24. Diekmann, M. *et al.* Patterns of long-term vegetation change vary between different types of semi-natural grasslands in Western and Central Europe. *J Veg Sci* **30**, 187–202 (2019).
25. Hundt, R. *Ökologisch-geobotanische Untersuchungen an den mitteldeutschen Wiesengesellschaften unter besonderer Berücksichtigung ihres Wasserhaushaltes und ihrer Veränderung durch die Intensivbewirtschaftung.* (Wehry-Druck OHG, 2001).
26. Kuhn, G., Heinz, S. & Mayer, F. Grünlandmonitoring Bayern. Ersterhebung der Vegetation 2002–2008. *Schriftenreihe LfL Bayerische Landesanstalt für Landwirtschaft* **3**, 1–161 (2011).
27. Raehse, S. *Veränderungen der hessischen Grünlandvegetation seit Beginn der 50er Jahre am Beispiel ausgewählter Tal- und Bergregionen Nord- und Mittelhessens.* (University Press GmbH, 2001).
28. Scheidel, U. & Bruelheide, H. Versuche zur Beweidung von Bergwiesen im Harz. *Hercynia N.F.* **37**, 87–101 (2004).
29. Sommer, S. & Hachmöller, B. Auswertung der Vegetationsaufnahmen von Dauerbeobachtungsflächen auf Bergwiesen im NSG Oelsen bei variiertem Mahd im Vergleich zur Brache. *Ber. Arbeitsgem. Sächs. Bot. N.F.* **18**, 99–135 (2001).
30. Wegener, U. Vegetationswandel des Berggrünlands nach Untersuchungen von 1954 bis 2016 - Wege zur Erhaltung der Bergwiesen. Mountain grasslands vegetation change after research from 1954 to 2016 - ways to preserve mountain meadows. *Abhandlungen und Berichte aus dem Museum Heineanum* **11**, 35–101 (2018).
31. Wittig, B., Müller, J. & Mahnke-Ritoff, A. Talauen-Glatthaferwiesen im Verdener Wesertal (Niedersachsen). *Tuexenia* **39**, 249–265 (2019).
32. Heinrich, W., Marstaller, R. & Voigt, W. Eine Langzeitstudie zur Sukzession in Halbtrockenrasen – Strukturwandlungen in einer Dauerbeobachtungsfläche im Naturschutzgebiet “Leutrat und Cospoth” bei Jena (Thüringen). *Artenschutzreport Jena* **30**, 1–80 (2012).
33. Hüllbusch, E., Brand, L. M., Ende, P. & Dengler, J. Little vegetation change during two decades in a dry grassland complex in the Biosphere Reserve Schorfheide-Chorin (NE Germany). *Tuexenia* **36**, 395–412 (2016).
34. Knapp, R. Dauerflächen-Untersuchungen über die Einwirkung von Haustieren und Wild während trockener und feuchter Zeiten in Mesobromion-Halbtrockenrasen in Hessen. *Mitt. Florist.-Soziol. Arbeitsgem. N.F.* **19/20**, 269–274 (1977).
35. Matesanz, S., Brooker, R. W., Valladares, F. & Klotz, S. Temporal dynamics of marginal steppic vegetation over a 26-year period of substantial environmental change: Temporal dynamics of marginal steppic vegetation over a 26-year period. *Journal of Vegetation Science* **20**, 299–310 (2009).
36. Schwabe, A., Zehm, A., Nobis, M., Storm, C. & Süß, K. Auswirkungen von Schaf-Erstbeweidung auf die Vegetation primär basenreicher Sand-Ökosysteme. *NNA Berichte* **1/2004**, 39–54 (2004).
37. Schwabe, A., Süß, K. & Storm, C. What are the long-term effects of livestock grazing in steppic sandy grassland with high conservation value? Results from a 12-year field study. *Tuexenia* **33**, 189–212 (2013).
38. Peppeler-Lisbach, C., Stanik, N., Könitz, N. & Rosenthal, G. Long-term vegetation changes in *Nardus* grasslands indicate eutrophication, recovery from acidification, and management change as the main drivers. *Appl Veg Sci* **23**, 508–521 (2020).
39. Peppeler-Lisbach, C. & Könitz, N. Vegetationsveränderungen in Borstgrasrasen des Werra-Meißner-Gebietes (Hessen, Niedersachsen) nach 25 Jahren. *Tuexenia* **37**, 201–228 (2017).
40. Wittig, B., Müller, J., Quast, R. & Miehllich, H. *Arnica montana* in Calluna-Heiden auf dem Schießplatz Unterlüß (Niedersachsen). *Tuexenia* **40**, 131–146 (2020).
41. Rumpf, S. B. *et al.* Range dynamics of mountain plants decrease with elevation. *Proc Natl Acad Sci USA* **115**, 1848–1853 (2018).
42. Kudernatsch, T. *et al.* Vegetationsveränderungen alpiner Kalk-Magerrasen im Nationalpark Berchtesgaden während der letzten drei Jahrzehnte. *Tuexenia* **36**, 205–221 (2016).
43. Poschlod, P. *et al.* Long-term monitoring in rivers of south Germany since the 1970ies. Macrophytes as indicators for the assessment of water quality. in *Long-term ecological research. Between Theory and Application* (eds. Müller, F., Baessler, C., Schubert, H. & Klotz, S.) 189–199 (Springer, 2006).
44. Dierschke, H. Dynamik und Konstanz an naturnahen Flussufern. 27 Jahre Dauerflächenuntersuchungen am Oderufer (Harzvorland). *Braunschweiger Geobotanische Arbeiten* **9**, 119–138 (2008).
45. Kreyling, J. *et al.* Rewetting does not return drained fen peatlands to their old selves. *Nat Commun* **12**, 5693 (2021).
46. Bohn, U. & Schniotalle, S. *Hochmoor-, Grünland- und Waldrenaturierung im Naturschutzgebiet ‘Rotes Moor’/Hohe Rhön 1981–2001.* (Landwirtschaftsverlag, 2008).
47. Koch, M. & Jurasinski, G. Four decades of vegetation development in a percolation mire complex following intensive drainage and abandonment. *Plant Ecology & Diversity* **8**, 49–60 (2015).
48. Walther, K. Die Vegetation des Maujahn 1984. Wiederholung der vegetationskundlichen Untersuchung eines wendländischen Moores. *Tuexenia* **6**, 145–193 (1986).
49. Berg, C. & Mahn, E.-G. Anthropogene Vegetationsveränderungen der Straßenrandvegetation in den letzten 30 Jahren - die Glatthaferwiesen des Raumes Halle/Saale. *Tuexenia* **10**, 185–195 (1990).
50. Meyer, S., Wesche, K., Krause, B. & Leuschner, C. Dramatic losses of specialist arable plants in Central Germany since the 1950s/60s - a cross-regional analysis. *Diversity Distribution* **19**, 1175–1187 (2013).
51. Meyer, S., Wesche, K., Krause, B. & Leuschner, C. Veränderungen in der Segetalflora in den letzten Jahrzehnten und mögliche Konsequenzen für Agrarvögel. *Julius-Kühn-Archiv* **442**, 64–78 (2013).
52. Kutzelnigg, H. Veränderungen der Ackerwildkrautflora im Gebiet um Moers/Niederrhein seit 1950 und ihre Ursachen. *Tuexenia* **4**, 81–102 (1984).

53. Milligan, G., Rose, R. J. & Marrs, R. H. Winners and losers in a long-term study of vegetation change at Moor House NNR: Effects of sheep-grazing and its removal on British upland vegetation. *Ecological Indicators* **68**, 89–101 (2016).
54. Wittig, B., Waldman, T. & Diekmann, M. Veränderungen der Grünlandvegetation im Holtumer Moor über vier Jahrzehnte. *Hercynia N.F.* **40**, 285–300 (2007).
55. Henning, K., Lorenz, A., von Oheimb, G., Härdtle, W. & Tischew, S. Year-round cattle and horse grazing supports the restoration of abandoned, dry sandy grassland and heathland communities by suppressing *Calamagrostis epigejos* and enhancing species richness. *Journal for Nature Conservation* **40**, 120–130 (2017).
56. Blüml, V. Langfristige Veränderungen von Flora und Vegetation des Grünlandes in der Dümmerniederung (Niedersachsen) unter dem Einfluss von Naturschutzmaßnahmen. (Bremen, 2011).
57. Von Oheimb, G. *et al.* *Halboffene Weidelandchaft Höltigbaum. Perspektiven für den Erhalt und die naturverträgliche Nutzung von Offenlandlebensräumen.* (Landwirtschaftsverlag, 2006).
58. Dornelas, M. *et al.* BioTIME: A database of biodiversity time series for the Anthropocene. *Global Ecol Biogeogr* **27**, 760–786 (2018).
59. Pimm, S. L. *et al.* The biodiversity of species and their rates of extinction, distribution, and protection. *Science* **344**, 1246752–1246752 (2014).
60. Barnosky, A. D. *et al.* Has the Earth's sixth mass extinction already arrived? *Nature* **471**, 51–57 (2011).
61. Vellend, M. The Biodiversity Conservation Paradox. *Am. Sci.* **105**, 94 (2017).
62. Cardinale, B. J., Gonzalez, A., Allington, G. R. H. & Loreau, M. Is local biodiversity declining or not? A summary of the debate over analysis of species richness time trends. *Biological Conservation* **219**, 175–183 (2018).
63. Perring, M. P. *et al.* Understanding context dependency in the response of forest understorey plant communities to nitrogen deposition. *Environmental Pollution* **242**, 1787–1799 (2018).
64. Steinbauer, M. J. *et al.* Accelerated increase in plant species richness on mountain summits is linked to warming. *Nature* **556**, 231–234 (2018).
65. Braun-Blanquet, J. Prinzipien einer Systematik der Pflanzengesellschaften auf floristischer Grundlage. *Jahrb. St. Gallischen Naturwiss. Ges.* **57**, 305–351 (1921).
66. Becking, R. W. The Zürich-Montpellier school of phytosociology. *Bot. Rev.* **23**, 411–488 (1957).
67. Bruehlheide, H. *et al.* sPlot – A new tool for global vegetation analyses. *J Veg Sci* **30**, 161–186 (2019).
68. O L Pescott, T A Humphrey & K J Walker. A short guide to using British and Irish plant occurrence data for research. <https://doi.org/10.13140/RG.2.2.33746.86720> (2018).
69. Eichenberg, D. *et al.* Widespread decline in Central European plant diversity across six decades. *Global Change Biology* **27**, 1097–1110 (2021).
70. Chytrý, M. *et al.* European Vegetation Archive (EVA): an integrated database of European vegetation plots. *Appl Veg Sci* **19**, 173–180 (2016).
71. Van der Maarel, E. Transformation of cover-abundance values in phytosociology and its effects on community similarity. *Vegetatio* **39**, 97–114 (1979).
72. Tichý, L. *et al.* Optimal transformation of species cover for vegetation classification. *Appl Veg Sci* **23**, 710–717 (2020).
73. Podani, J. Braun-Blanquet's legacy and data analysis in vegetation science. *Journal of Vegetation Science* **17**, 113–117 (2006).
74. Londo, G. Dezimalskala für die vegetationskundliche Aufnahme von Dauerquadraten. in *Sukzessionsforschung* (ed. Schmidt, W.). *Ber. Int. Symp. Int. Vereinig. Vegetationsk. Rinteln* vol. 1973, 613–617 (Cramer, 1975).
75. Bruehlheide, H. & Luginbühl, U. Peeking at ecosystem stability: making use of a natural disturbance experiment to analyze resistance and resilience. *Ecology* **90**, 1314–1325 (2009).
76. Hennekens, S. M. & Schaminée, J. H. J. TURBOVEG, a comprehensive data base management system for vegetation data. *J. Veg. Sc.* **12**, 589–591 (2001).
77. Gaston, K. J. & Curnutt, J. L. The dynamics of abundance-range size relationships. *Oikos* **81**, 38 (1998).
78. Gaston, K. J. *et al.* Abundance-occupancy relationships. *J Appl Ecology* **37**, 39–59 (2000).
79. Sporbert, M. *et al.* Testing macroecological abundance patterns: The relationship between local abundance and range size, range position and climatic suitability among European vascular plants. *J Biogeogr* jbi.13926, <https://doi.org/10.1111/jbi.13926> (2020).
80. European Commission. *Report on the Conservation Status of Habitat Types and Species as required under Article 17 of the Habitats Directive.* <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:52009DC0358> (2009).
81. Poschlod, P. *Geschichte der Kulturlandschaft.* (Ulmer, 2017).
82. McGill, B., Enquist, B., Weiher, E. & Westoby, M. Rebuilding community ecology from functional traits. *Trends in Ecology & Evolution* **21**, 178–185 (2006).
83. Jandt, U. *et al.* More losses than gains during one century of plant biodiversity change in Germany. *Nature* <https://doi.org/10.1038/s41586-022-05320-w> (2022).
84. Schaminée, J. H. J., Hennekens, S. M., Chytrý, M. & Rodwell, J. S. Vegetation-plot data and databases in Europe: an overview. *Preslia* **81**, 173–185 (2009).
85. ESA. Land Cover CCI product user guide ver. 2. Tech. Rep. https://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf (2017).
86. Kadmon, R., Farber, O. & Danin, A. Effect of roadside bias on the accuracy of predictive maps produced by bioclimatic models. *Ecological Applications* **14**, 401–413 (2004).
87. Davies, C. E., Moss, D. & Hill, M. O. *EUNIS Habitat Classification Revised 2004.* 310 <https://www.eea.europa.eu/data-and-maps/data/eunis-habitat-classification/documentation/eunis-2004-report.pdf/download> (2004).
88. Chytrý, M. *et al.* EUNIS Habitat Classification: Expert system, characteristic species combinations and distribution maps of European habitats. *Appl Veg Sci* **23**, 648–675 (2020).
89. Bruehlheide, H., Tichý, L., Chytrý, M. & Jansen, F. Implementing the formal language of the vegetation classification expert systems (ESy) in the statistical computing environment R. *Appl Veg Sci*, <https://doi.org/10.1111/avsc.12562> (2021).
90. Jandt, U., Bruehlheide, H. & ReSurveyGermany Consortium. ReSurvey Germany: vegetation-plot resurvey data from Germany. *German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig* <https://doi.org/10.25829/idiv.3514-0qsq70> (2022).
91. Jansen, F. & Dengler, J. GermanSL - eine universelle taxonomische Referenzliste für Vegetationsdatenbanken. *Tuexenia* **28**, 239–253 (2008).
92. Wisskirchen, R. & Haeupler, H. *Standardliste der Farn- und Blütenpflanzen Deutschlands.* (Ulmer, 1998).
93. Jansen, F. & Dengler, J. Plant names in vegetation databases—a neglected source of bias. *Journal of Vegetation Science* **21**, 1179–1186 (2010).
94. Fischer, H. S. On the combination of species cover values from different vegetation layers. *Applied Vegetation Science* **18**, 169–170 (2015).
95. Schwabe, A. & Kratochwil, A. Pflanzensoziologische Dauerflächen-Untersuchungen im Bannwald 'Flüh' (Südschwarzwald) unter besonderer Berücksichtigung der Weidfeld-Sukzession. *Standort. Wald* **49**, 5–49 (2015).
96. Poschlod, P., Schreiber, K.-F., Mitlacher, K., Römermann, C. & Bernhardt-Römermann, M. Entwicklung der Vegetation und ihre naturschutzfachliche Bewertung. in *Landschaftspflege und Naturschutz im Extensivgrünland. 30 Jahre Offenhaltungsversuche Baden-Württemberg* (eds. Schreiber, K.-F., Brauckmann, H.-J., Broll, G., Krebs, S. & Poschlod, P.) vol. 97 243–288 (2009).
97. Gonzalez, A. *et al.* Estimating local biodiversity change: a critique of papers claiming no net loss of local diversity. *Ecology* **97**, 1949–1960 (2016).

Acknowledgements

We are grateful to surveyors who recorded vegetation in the field and provided these data. We acknowledge those data contributors who made their data available to us or helped in recording these data: Thea Dittmann, Alexandra Erfmeier, Bernd Gerken, Kerstin Günther, Sabine Heinz, Wilfried Hakes, Heike Heklau, Alfons Henrichfreise, Elisabeth Hüllbusch, Andreas Huwer, Anneke Immoor, Sophie Luise Kühn, Benjamin Krause, Sebastian Leonhardt, Thomas Meineke, Jutta Rach, Jennifer Reinecke, Ulrich Scheidel and Immo Vollmer. We thank Diana Bowler and Netra Bhandari for the analysis of spatial representativeness. The assistance of the iDiv Data & Code Unit (Anahita Kazem, Ludmilla Figueiredo) for archiving the dataset is greatly acknowledged. The manuscript very much benefitted from the comments of Javier Gamarra and two other anonymous reviewers. We very much appreciate the support for the strategic project sMon by the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German Research Foundation (DFG-FZT 118, 202548816).

Author contributions

U.J. and H.B. conceived the idea for the project and compiled the data. All authors were involved in collecting datasets, which were harmonized by U.J. and H.B. The first draft, the figures and tables were produced by H.B. All authors discussed and commented on the manuscript and contributed to the revised versions.

Funding

Open Access funding enabled and organized by Projekt DEAL.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41597-022-01688-6>.

Correspondence and requests for materials should be addressed to H.B.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022

Ute Jandt ^{1,2,48}, Helge Bruelheide ^{1,2,48} ✉, Christian Berg ³, Markus Bernhardt-Römermann ^{2,4}, Volker Blüml ⁵, Frank Bode⁶, Jürgen Dengler ^{2,7,8}, Martin Diekmann⁹, Hartmut Dierschke¹⁰, Inken Doerfler¹¹, Ute Döring¹², Stefan Dullinger ¹³, Werner Härdtle ¹⁴, Sylvia Haider^{1,2}, Thilo Heinken ¹⁵, Peter Horchler¹⁶, Florian Jansen ¹⁷, Thomas Kudernatsch¹⁸, Gisbert Kuhn ¹⁹, Martin Lindner^{2,20}, Silvia Matesanz²¹, Katrin Metzger²², Stefan Meyer²³, Frank Müller ²⁴, Norbert Müller ²⁵, Tobias Naaf ²⁶, Cord Peppeler-Lisbach ²⁷, Peter Poschold²⁸, Christiane Roscher^{2,29}, Gert Rosenthal³⁰, Sabine B. Rumpf ^{13,31}, Wolfgang Schmidt ³², Joachim Schrautzer³³, Angelika Schwabe ³⁴, Peter Schwartz³⁵, Thomas Sperle³⁶, Nils Stanik ³⁰, Hans-Georg Stroh³⁷, Christian Storm³⁸, Winfried Voigt³⁹, Andreas von Heßberg⁴⁰, Goddert von Oheimb ⁴¹, Eva-Rosa Wagner⁴², Uwe Wegener⁴³, Karsten Wesche ^{2,44,45}, Burghard Wittig ^{9,46} & Monika Wulf ^{26,47}

¹Institute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108, Halle, Germany. ²German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Pusckstr. 4, 04103, Leipzig, Germany. ³Karl-Franzens-Universität Graz, Institute for Biology, Holteigasse 6, 8010, Graz, Austria. ⁴Institute of Ecology and Evolution, Friedrich Schiller University Jena, Dornburger Str. 159, 07743, Jena, Germany. ⁵BMS - Umweltplanung, Freiheitsweg 38A, 49086, Osnabrück, Germany. ⁶Abteilung Forschungsförderung, Karlsruher Institut für Technologie (KIT), Kaiserstraße 12, 76131, Karlsruhe, Germany. ⁷Vegetation Ecology Group, Institute of Natural Resource Sciences (IUNR), Zurich University of Applied Sciences (ZHAW), Grüentalstr. 14, 8820, Wädenswil, Switzerland. ⁸Plant Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER),

Universitätsstr. 30, Bayreuth, 95447, Germany. ⁹Vegetation Ecology and Conservation Biology, Institute of Ecology, FB 2, University of Bremen, Bremen, Germany. ¹⁰Vegetation Analysis and Phytodiversity, Albrecht-von-Haller-Institute of Plant Sciences, Georg-August-University of Göttingen, Untere Karspüle 2, D-37073, Göttingen, Germany. ¹¹Vegetation Science and Nature Conservation Group, Institute for Biology and Environmental Sciences, University of Oldenburg, 2611, Oldenburg, Germany. ¹²Auf der Wessel 47, 37085, Göttingen, Germany. ¹³Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, 1030, Vienna, Austria. ¹⁴Leuphana University of Lüneburg, Institute of Ecology, Universitätsallee 1, 21335, Lüneburg, Germany. ¹⁵General Botany, Institute of Biochemistry and Biology, University of Potsdam, Maulbeerallee 3, 14469, Potsdam, Germany. ¹⁶Federal Institute of Hydrology, Department Vegetation Studies, Landscape Management, Am Mainzer Tor 1, 56068, Koblenz, Germany. ¹⁷Faculty of Agricultural and Environmental Sciences, Rostock University, Justus-von-Liebig-Weg 6, 18059, Rostock, Germany. ¹⁸Bavarian State Institute of Forestry, Hans-Carl-von-Carlowitz-Platz 1, 85354, Freising, Germany. ¹⁹Institut für Agrarökologie und Biologischen Landbau, AG Vegetationsökologie und -monitoring, Bayerische Landesanstalt für Landwirtschaft, Lange Point 12, 85354, Freising, Germany. ²⁰Institute of Biology/Biology Education, Martin Luther University Halle-Wittenberg, Weinbergweg 10, 06120, Halle, Germany. ²¹Universidad Rey Juan Carlos, Area de Biodiversidad y Conservación, Móstoles, Madrid, 28933, Spain. ²²Ministerium für Wissenschaft, Energie, Klimaschutz und Umwelt des Landes Sachsen-Anhalt, Leipziger Straße 58, 39112, Magdeburg, Germany. ²³Plant Ecology and Ecosystems Research, Albrecht von Haller Institute of Plant Sciences, University of Göttingen, Untere Karspüle 2, 37073, Göttingen, Germany. ²⁴Institute of Botany, TU Dresden, Mommsenstr. 13, 01062, Dresden, Germany. ²⁵Dep. Landscape Management & Restoration Ecology, Fachhochschule Erfurt, Leipzigerstr. 77, 99085, Erfurt, Germany. ²⁶Leibniz Centre for Agricultural Landscape Research (ZALF), Eberswalder Straße 84, 15374, Müncheberg, Germany. ²⁷Landscape Ecology Group, Institute for Biology and Environmental Sciences, University of Oldenburg, Carl von Ossietzky Str. 9–11, 26129, Oldenburg, Germany. ²⁸Ecology and Conservation Biology, Institute of Plant Sciences, Faculty of Biology and Preclinical Medicine, University of Regensburg, Universitaetsstrasse 31, 93053, Regensburg, Germany. ²⁹Department of Physiological Diversity, UFZ, Helmholtz Centre for Environmental Research, Puschstr. 4, 04103, Leipzig, Germany. ³⁰Department of Landscape and Vegetation Ecology, University of Kassel, Gottschalkstrasse 26a, 34127, Kassel, Germany. ³¹University of Basel, Department of Environmental Sciences, Bernoullistrasse 32, 4056, Basel, Switzerland. ³²Department of Silviculture and Forest Ecology of the Temperate Zones, Georg-August-University Göttingen, Büsgenweg 1, D-37077, Göttingen, Germany. ³³Institute for Ecosystem Research, Kiel University, Olshausenstraße 75, 24118, Kiel, Germany. ³⁴Faculty of Biology, Technical University Darmstadt, Schnittpahnstraße 4, 64287, Darmstadt, Germany. ³⁵Biologische Station Kreis Steinfurt e.V., Bahnhofstraße 71, 49545, Tecklenburg, Germany. ³⁶Vogtsstr. 3, 79183, Waldkirch, Germany. ³⁷büro áchero Vegetation and Environmental Consulting, Friedländer Straße 17a, 37133, Friedland, Germany. ³⁸Fachgebiet Chemische Pflanzenökologie, Fachbereich Biologie, Technische Universität Darmstadt, Schnittpahnstr. 10, D-64287, Darmstadt, Germany. ³⁹Institute of Ecology and Evolution, University of Jena, Dornburger Str. 159, 07743, Jena, Germany. ⁴⁰Bayceer - Uni Bayreuth, Universitätsstr. 30, 95447, Bayreuth, Germany. ⁴¹Technische Universität Dresden, Institute of General Ecology and Environmental Protection, Piener Straße 7, 01737, Tharandt, Germany. ⁴²Engbertsheide 9, 49324, Melle, Germany. ⁴³Meisenweg 27, 38820, Halberstadt, Germany. ⁴⁴Botany Department, Senckenberg Museum of Natural History Görlitz, Am Museum 1, 02826, Görlitz, Germany. ⁴⁵International Institute Zittau, Technische Universität Dresden, Markt 23, 02763, Zittau, Germany. ⁴⁶Lower Saxony Water Management, Coastal Protection and Nature Conservation Agency, Betriebsstelle Lüneburg, Standort Verden, Bürgermeister Münchmeyer Str. 6, 27283, Verden, Germany. ⁴⁷Institute of Biochemistry and Biology, University of Potsdam, Maulbeerallee 3, 14469, Potsdam, Germany. ⁴⁸These authors contributed equally: Ute Jandt, Helge Bruelheide. ✉e-mail: helge.bruehlheide@botanik.uni-halle.de